

# ATP-independent Renaturation of Complementary DNA Strands by the Mutant *recA1* Protein from *Escherichia coli*\*

(Received for publication, March 10, 1986)

Floyd R. Bryant‡ and I. R. Lehman

From the Department of Biochemistry, Stanford University School of Medicine, Stanford, California 94305

In an effort to clarify the requirement for ATP in the *recA* protein-promoted renaturation of complementary DNA strands, we have analyzed the mutant *recA1* protein which lacks single-stranded DNA-dependent ATPase activity at pH 7.5. Like the wild type, the *recA1* protein binds to single-stranded DNA with a stoichiometry of one monomer per approximately four nucleotides. However, unlike the wild type, the mutant protein is dissociated from single-stranded DNA in the presence of ATP or ADP. The ATP analogue adenosine 5'-O-3'-(thiotriphosphate) appears to stabilize the binding of *recA1* protein to single-stranded DNA but does not elicit the stoichiometry of 1 monomer/8 nucleotides or the formation of highly condensed protein-DNA networks that are characteristic of the wild type *recA* protein in the presence of this analogue. The *recA1* protein does not catalyze DNA renaturation in the presence of ATP, consistent with the dissociation of *recA1* protein from single-stranded DNA under these conditions. However, it does promote a pattern of Mg<sup>2+</sup>-dependent renaturation identical to that found for wild type *recA* protein.

The renaturation of complementary DNA strands catalyzed by the *recA* protein of *Escherichia coli* is stimulated by ATP (1). However, under certain conditions, renaturation can proceed efficiently in the absence of a nucleotide cofactor (2). In an effort to clarify the role of ATP in the *recA* protein-catalyzed renaturation of DNA, we have compared the wild type *recA* protein with the mutant *recA1* protein that lacks a ssDNA<sup>1</sup>-dependent ATPase (3). We have found that the *recA1* protein does not catalyze renaturation in the presence of ATP. However, it does promote Mg<sup>2+</sup>-dependent renaturation in a manner identical to that of the wild type *recA* protein. Thus, only ATP-dependent DNA renaturation is defective in the *recA1* mutant.

## EXPERIMENTAL PROCEDURES

### Materials

*RecA* protein was purified to homogeneity as described (4). *RecA1* protein was generously provided by Dr. Steve West (Yale University). S1 nuclease, calf thymus DNA, and phosphocreatine were from

\* This work was supported by Grant GM06196 from the National Institutes of Health. The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

‡ Supported by Postdoctoral Fellowship DRG579 from the Damon Runyon-Walter Winchell Cancer Fund and the Bank of America-Giannini Foundation. Present address: Dept. of Biochemistry, Johns Hopkins University School of Public Health, Baltimore, MD 21205.

<sup>1</sup> The abbreviations used are: ssDNA, single-stranded DNA;  $\phi$ X, bacteriophage  $\phi$ X174; ATP $\gamma$ S, adenosine 5'-O-3'-(thiotriphosphate).

Sigma. DNaseI and snake venom phosphodiesterase were from Worthington. ATP $\gamma$ S and creatine phosphokinase were from Boehringer Mannheim. GF/C filters were from Whatman and nitrocellulose filters (HAWP, 0.45- $\mu$ m pore diameter) were from Millipore. [<sup>3</sup>H]ATP, [ $\gamma$ -<sup>32</sup>P]ATP, and [ $\gamma$ -<sup>35</sup>S]ATP $\gamma$ S were from Amersham.

Unlabeled and <sup>3</sup>H-labeled circular  $\phi$ X ssDNA ((+) strand) and <sup>3</sup>H-labeled linear duplex  $\phi$ X DNA were prepared as described (5). <sup>3</sup>H-labeled linear duplex  $\phi$ X DNA was denatured as described (2).

### Methods

**Nuclease Protection Assay**—Nuclease protection assays were carried out as described previously (6). Reaction mixtures (500  $\mu$ l) contained 25 mM Tris·HCl (pH 7.5), 10 mM MgCl<sub>2</sub>, 30  $\mu$ M [<sup>3</sup>H] $\phi$ X ssDNA, 5% glycerol, 1  $\mu$ M *recA* or *recA1* protein, and the indicated nucleotide cofactors. When an ATP regeneration system was included, it consisted of 8 mM phosphocreatine and 40  $\mu$ g/ml creatine phosphokinase. After a 5-min incubation at 25 °C, DNaseI (10  $\mu$ g) and venom phosphodiesterase (10  $\mu$ g) were added, and the amount of [<sup>3</sup>H]ssDNA remaining resistant to nuclease digestion was determined as previously described (6).

**Nitrocellulose Filter-binding Assay**—All nitrocellulose filter-binding assays were carried out in solutions containing 25 mM Tris·HCl (pH 7.5), 10 mM MgCl<sub>2</sub>, 5% glycerol, and the indicated amounts of  $\phi$ X ssDNA, nucleotide cofactor, and *recA* or *recA1* protein unless otherwise indicated. Following incubation, the reaction mixtures were filtered on KOH-treated nitrocellulose filters, washed with 1 ml of reaction buffer, dried, and then assayed for the appropriate radioactivity. All procedures were carried out at 25 °C.

***RecA* Protein Transfer Kinetics**—In the standard transfer reaction, *recA* or *recA1* protein (1  $\mu$ M) was preincubated with 30  $\mu$ M [<sup>3</sup>H] $\phi$ X ssDNA in 25 mM Tris·HCl (pH 7.5), 10 mM MgCl<sub>2</sub>, and 1 mM dithiothreitol for 5 min at 25 °C. The preincubation solution was then mixed with an equal volume of challenge solution containing 25 mM Tris·HCl (pH 7.5), 10 mM MgCl<sub>2</sub>, 1 mM dithiothreitol, and 150  $\mu$ g/ml of heat-denatured calf thymus DNA; the mixture was kept at 25 °C. At various times, aliquots (75  $\mu$ l) were removed and assayed by nitrocellulose filter binding as described above. Alternatively, aliquots (75  $\mu$ l) were quenched with ATP $\gamma$ S (500  $\mu$ M) and assayed by nuclease digestion as described previously (6). In the nitrocellulose filter assay, zero time points were measured by mixing 37.5  $\mu$ l of the preincubation solution with 37.5  $\mu$ l of reaction buffer, followed by filtering. Zero time points for the nuclease protection assay were measured as previously described (6).

**DNA Renaturation Assay**—Renaturation assays were carried out as described (2). Reaction mixtures contained 25 mM Tris·HCl (pH 7.5), 5% glycerol, <sup>3</sup>H-labeled denatured *Pst*I cleaved  $\phi$ X DNA, MgCl<sub>2</sub>, nucleotides, and *recA* or *recA1* protein at the concentrations given in the figure legends.

**ssDNA-dependent ATPase Assay**—Steady-state ATP hydrolysis catalyzed by *recA* protein was measured using a thin-layer chromatography method that has been previously described (1). Pre-steady-state hydrolysis was measured using the following Norit adsorption assay (7). *RecA* or *recA1* protein (12  $\mu$ M) was preincubated with 100  $\mu$ M  $\phi$ X ssDNA in 25 mM Tris·HCl (pH 7.5), 10 mM MgCl<sub>2</sub>, 1 mM dithiothreitol, and 5% glycerol at 25 °C for 5 min. The preincubation solution (100  $\mu$ l) was then mixed with an equal volume of solution containing 25 mM Tris·HCl (pH 7.5), 10 mM MgCl<sub>2</sub>, 1 mM dithiothreitol, 5% glycerol, and 400  $\mu$ M [ $\gamma$ -<sup>32</sup>P]ATP (specific activity approximately 70,000 cpm/nmol) and kept at 25 °C. After various time intervals, the reaction mixtures were quenched with 200  $\mu$ l of HCl (0.5 N) and kept on ice. The quenched solutions were then mixed

with 1 ml of acid-washed Norit (30 mg/ml in 0.1 N HCl), vortexed, and then centrifuged. The supernatants (1.0 ml) were mixed with 6 ml of aqueous scintillation-counting solution and then assayed for  $^{32}\text{P}$ . Zero time points were measured by adding 100  $\mu\text{l}$  of the preincubation solution first to 0.2 ml HCl (0.1 N) followed by the addition of 100  $\mu\text{l}$  of the  $[\gamma\text{-}^{32}\text{P}]\text{ATP}$ -containing solution.

## RESULTS

**Binding of *recA* and *recA1* Proteins to ssDNA: Nuclease Protection.**—We have described a nuclease digestion protection method to quantitate the binding of *recA* monomers to ssDNA (6). The stoichiometries of protection obtained using this assay are summarized in Table I. In the absence of nucleotide, each *recA* monomer protected approximately four nucleotides of ssDNA from digestion. The same stoichiometry was found for complexes formed in the presence of ATP (plus an ATP regeneration system). In the presence of ATP without the ATP regeneration system or in the presence of ADP, all of the DNA remained susceptible to nuclease digestion, presumably because of the weaker binding of *recA* protein to ssDNA under these conditions. In the presence of ATP $\gamma\text{S}$ , a nonhydrolyzable analog of ATP, each *recA* monomer protected approximately eight nucleotides of ssDNA, twice that seen in the absence of ATP $\gamma\text{S}$ .

The binding of the mutant *recA1* protein to ssDNA is also summarized in Table I. In the absence of nucleotide, the *recA1* protein showed a stoichiometry similar to the wild type protein, with each *recA1* monomer protecting approximately 3.5 nucleotides of ssDNA from digestion. In the presence of ATP (plus or minus an ATP regeneration system) or ADP, there was no protection by *recA1* protein. In the presence of ATP $\gamma\text{S}$ , the stoichiometry of protection remained at 1 *recA1* monomer/4.0 nucleotides rather than increasing 2-fold as it did in the case of the wild type *recA* protein.

**Binding of *recA* and *recA1* Proteins to ssDNA: Nitrocellulose Filter Binding.**—The binding of *recA* and *recA1* proteins to ssDNA was also examined by nitrocellulose filter binding. A fixed concentration of  $[\text{H}^3]\phi\text{X}$  ssDNA (30  $\mu\text{M}$ ) was incubated with increasing concentrations of *recA* or *recA1* protein and the amount of ssDNA retained on nitrocellulose filters was measured. As shown in Fig. 1, the *recA* and *recA1* proteins retained ssDNA with a similar efficiency. For the *recA* protein, approximately 1 *recA* monomer/30 nucleotides of ssDNA was required to retain >80% of the ssDNA molecules, corresponding to approximately 200 *recA* monomers for every ssDNA molecule. Since, in principle, the binding of a single *recA* monomer to each ssDNA molecule should be sufficient for retention, the value of 200 probably reflects a cooperative mode of binding in which subsaturating levels of *recA* protein bind to ssDNA as clusters of contiguous monomers. The binding of *recA1* protein to ssDNA showed a similar behavior.

The nitrocellulose filter-binding assay was used to evaluate




FIG. 1. Retention of  $\phi\text{X}$  ssDNA on nitrocellulose filters by *recA* and *recA1* proteins: dependence on protein concentration. Nitrocellulose filter binding was carried out as described under "Experimental Procedures." Reaction solutions contained 30  $\mu\text{M}$   $[\text{H}^3]\phi\text{X}$  ssDNA and the indicated concentrations of protein. (●), *recA* protein; (○), *recA1* protein. The points represent the percentage of the total DNA retained on a nitrocellulose filter.

the effects of various nucleotides on the binding of *recA* and *recA1* proteins to ssDNA.  $[\text{H}^3]\phi\text{X}$  ssDNA (30  $\mu\text{M}$ ) was incubated with sufficient concentrations of *recA* or *recA1* protein (1  $\mu\text{M}$ ) to give approximately 80% retention on nitrocellulose filters as judged by the titration curves above. Various nucleotides were then added to the mixture, and the efficiency of retention of the  $[\text{H}^3]\phi\text{X}$  ssDNA on a nitrocellulose filter was measured (Table II). No ssDNA was retained by either *recA* or *recA1* protein in the absence of  $\text{MgCl}_2$ . In the presence of 10 mM  $\text{MgCl}_2$ , both *recA* and *recA1* protein gave the expected retention. Treatment of the *recA*·ssDNA complex with ATP, ADP, or ATP $\gamma\text{S}$  resulted in no loss of efficiency of retention. In contrast, when the *recA1*·ssDNA complex was treated with either ATP or ADP, an almost complete loss of binding was observed; ATP $\gamma\text{S}$  did not decrease the filter binding efficiency. These results are consistent with the nuclease protection results in the previous section and indicate that ATP and ADP promote the rapid dissociation of *recA1* protein from ssDNA. Interestingly, dTTP caused a complete loss of binding for both the *recA* and *recA1* proteins.

**Transfer of *recA* and *recA1* Proteins between ssDNA Molecules.**—The nuclease protection assay can be used to measure the transfer of *recA* protein from one ssDNA molecule to another and to determine the effects of various nucleotides on this process (6). In a typical experiment (Fig. 2), *recA* protein (1  $\mu\text{M}$ ) was incubated with  $[\text{H}^3]\phi\text{X}$  ssDNA (30  $\mu\text{M}$ ) to form a *recA*·ssDNA complex. The transfer reaction was initiated by the addition of unlabeled challenge DNA (15 eq) to the preformed complexes. The reactions were then quenched after various times intervals by the addition of ATP $\gamma\text{S}$  which prevents dissociation of the *recA*·ssDNA complex. The complexes were then subjected to nuclease digestion. As shown in Fig. 2, there was a first order decay in the amount of  $[\text{H}^3]\phi\text{X}$  ssDNA protected from nuclease digestion as a function of time after the initiation of transfer, reflecting equilibration of *recA* protein between the labeled and unlabeled DNA populations. The  $t_{1/2}$  for equilibration using 15 eq of challenge DNA was approximately 5 min. A detailed study of *recA* protein transfer using this assay is found in Ref. 6.

The nuclease protection assay could not be used to measure

TABLE I

### Stoichiometry of protection of ssDNA by *recA* and *recA1* proteins

Protection stoichiometry measurements were carried out as described under "Experimental Procedures." Reaction solutions contained 30  $\mu\text{M}$   $[\text{H}^3]\phi\text{X}$  ssDNA, 1  $\mu\text{M}$  *recA* or *recA1* protein, and the indicated nucleotide cofactors (500  $\mu\text{M}$ ).

Nucleotide added	Nucleotides/ <i>recA</i> monomer	Nucleotides/ <i>recA1</i> monomer
None	4.2	3.5
ATP $\gamma\text{S}$	8.5	4.0
ATP (regeneration system added)	4.2	0
ATP (no regeneration system)	0	0
ADP	0	0




FIG. 2. *RecA* protein transfer: comparison of nuclease protection and nitrocellulose filter-binding assays. Reactions were carried out as described under "Experimental Procedures." Reaction mixtures contained  $30 \mu\text{M}$  [ $^3\text{H}$ ] $\phi\text{X}$  ssDNA and  $1 \mu\text{M}$  *recA* protein. Following a 5 min incubation, the transfer reactions were initiated by the addition of an equal volume of solution containing  $450 \mu\text{M}$  (15 eq) of heat-denatured salmon sperm DNA. At the times indicated, aliquots were quenched with ATP $\gamma\text{S}$  and assayed by: (O), the nuclease protection assay; (●), nitrocellulose filter binding (similar time courses were obtained with the filter-binding assay when the ATP $\gamma\text{S}$  quench was omitted). The points (O) represent the relative amount of the *recA* protein bound to the [ $^3\text{H}$ ]ssDNA; a value of 100 indicates that the *recA* protein is bound entirely to the [ $^3\text{H}$ ]DNA whereas a value of 0 indicates that the *recA* protein is completely equilibrated between the labeled and challenge ssDNA. The points (●) represent the relative amount of [ $^3\text{H}$ ]ssDNA retained on the filters; a value of 100 is equivalent to the amount of [ $^3\text{H}$ ]ssDNA retained by the *recA* protein in the absence of challenge DNA and a value of 0 is equivalent to the amount of [ $^3\text{H}$ ]ssDNA retained when *recA* protein ( $1 \mu\text{M}$ ) is added to a mixture of [ $^3\text{H}$ ]ssDNA ( $30 \mu\text{M}$ ) and challenge DNA ( $450 \mu\text{M}$ ).




FIG. 3. *RecA1* protein transfer: nitrocellulose filter-binding assay. Reactions were carried out as described under "Experimental Procedures." Reaction mixtures contained  $30 \mu\text{M}$  [ $^3\text{H}$ ] $\phi\text{X}$  ssDNA and  $1 \mu\text{M}$  *recA1* protein. (O), no nucleotide; (●),  $500 \mu\text{M}$  ATP $\gamma\text{S}$ . Following a 5 min incubation, the transfer reactions were initiated by the addition of an equal volume of solution containing  $450 \mu\text{M}$  (15 eq) of heat-denatured salmon sperm DNA. Aliquots were removed at the times indicated and assayed directly by nitrocellulose filter binding. The points represent the relative amount of [ $^3\text{H}$ ]ssDNA retained on the filters; a value of 100 is equivalent to the amount of [ $^3\text{H}$ ]ssDNA retained by the *recA1* protein in the absence of challenge DNA and a value of 0 is equivalent to the amount of [ $^3\text{H}$ ]ssDNA retained when *recA1* ( $1 \mu\text{M}$ ) is added to a mixture of [ $^3\text{H}$ ]ssDNA ( $30 \mu\text{M}$ ) and challenge DNA ( $450 \mu\text{M}$ ).

the transfer of *recA1* protein because of the relative instability of the *recA1*·ATP $\gamma\text{S}$  complex.<sup>2</sup> Under certain conditions it is, however, possible to measure *recA* protein transfer kinetics by nitrocellulose filter binding. A fixed concentration of wild type *recA* protein ( $1 \mu\text{M}$ ) was incubated with [ $^3\text{H}$ ] $\phi\text{X}$  ssDNA ( $30 \mu\text{M}$ ) to form a *recA*·ssDNA complex. As determined above, this level of *recA* protein is sufficient to cause retention of approximately 80% of the [ $^3\text{H}$ ] $\phi\text{X}$  ssDNA molecules. Transfer was initiated by the addition of unlabeled challenge DNA (15 eq) to the preformed complexes. At various times, aliquots were removed and filtered on nitrocellulose. As shown in Fig. 2, there was a first order decay in the retention of the [ $^3\text{H}$ ] $\phi\text{X}$  ssDNA as a function of time after the initiation of transfer. The time course of *recA* protein transfer obtained by the nitrocellulose filter-binding method was very similar to that obtained by nuclease protection and gave a  $t_{1/2}$  for equilibration of approximately 5 min.

The nitrocellulose filter-binding assay was also used to measure the transfer of *recA1* protein. As shown in Fig. 3, the  $t_{1/2}$  for equilibration of *recA1* protein with 15 eq of challenge DNA was approximately 1 min, about five times faster than for the wild type protein. Inclusion of ATP $\gamma\text{S}$  in the reaction mixture substantially reduced the rate of *recA1* protein transfer ( $t_{1/2} > 30$  min). ATP $\gamma\text{S}$  has a similar effect on the rate of transfer of the wild type *recA* protein (6).

**Formation of *recA1*·ssDNA·ATP $\gamma\text{S}$  Complexes**—The observation that ATP $\gamma\text{S}$  caused a substantial reduction in the rate of *recA1* protein transfer suggests that ATP $\gamma\text{S}$  does interact with the *recA1* protein, increasing its affinity for ssDNA. However, *recA1*·ssDNA·ATP $\gamma\text{S}$  complexes are different from *recA*·ssDNA·ATP $\gamma\text{S}$  complexes in several ways. First, as described above, ATP $\gamma\text{S}$  did not cause an increase in protection stoichiometry of ssDNA binding to the *recA1*

TABLE II

Retention of  $\phi\text{X}$  ssDNA on nitrocellulose filters by *recA* and *recA1* proteins

Condition	DNA retained	
	<i>recA</i>	<i>recA1</i>
	%	
No nucleotide	100	100
ATP	106	1
ADP	87	3
ATP $\gamma\text{S}$	100	98
dTTP	2	2
EDTA	1	1
ATP $\gamma\text{S}$ + EDTA <sup>a</sup>	100	5
-MgCl <sub>2</sub>	1	1

<sup>a</sup> The reaction mixture (+ATP $\gamma\text{S}$ ) was incubated for 5 min in the absence of EDTA. EDTA was then added to a concentration of 0.5 mM, and the incubation was continued for 5 min more before filtering.

protein. Second, electron microscopic studies showed that the addition of ATP $\gamma\text{S}$  to  $\phi\text{X}$  ssDNA ( $30 \mu\text{M}$ ) complexed with subsaturating levels of *recA1* protein ( $1 \mu\text{M}$ ) did not produce the highly condensed networks of protein and DNA, as it did in the case of the wild type *recA* protein.<sup>2</sup> Third, *recA*·ssDNA·ATP $\gamma\text{S}$  complexes were stable to the addition of EDTA whereas *recA1*·ssDNA·ATP $\gamma\text{S}$  complexes were completely disrupted, as judged by nitrocellulose filter binding (Table II).

In order to clarify the effect of ATP $\gamma\text{S}$  on the *recA1* protein, we examined directly the binding of ATP $\gamma\text{S}$  to *recA*·ssDNA and *recA1*·ssDNA complexes.  $\phi\text{X}$  ssDNA ( $30 \mu\text{M}$ ) was incubated with either *recA* or *recA1* protein ( $1 \mu\text{M}$ ) to form the desired protein·DNA complexes. [ $^{35}\text{S}$ ]ATP $\gamma\text{S}$  ( $200 \mu\text{M}$ ) was then added to the preformed complexes. The mixtures were filtered through nitrocellulose and the  $^{35}\text{S}$  retained on the filter measured.  $1.0 (\pm 0.05)$  molecule of [ $^{35}\text{S}$ ]ATP $\gamma\text{S}$  was

<sup>2</sup> F. R. Bryant and I. R. Lehman, unpublished observations.

bound for every recA monomer in the recA·ssDNA complex. A similar experiment with recA1 protein showed that approximately 0.8 molecule of [<sup>35</sup>S]ATP<sub>γ</sub>S was retained per monomer, indicating that the recA1 protein also binds a single equivalent of ATP<sub>γ</sub>S. However, the value for recA protein decreased with successive washings of the filter, suggesting that the recA1·ssDNA·ATP<sub>γ</sub>S complexes are subject to dissociation. In contrast, the recA·ssDNA·ATP<sub>γ</sub>S complexes were stable to multiple washings. Neither the recA nor recA1 protein retained [<sup>35</sup>S]ATP<sub>γ</sub>S on the nitrocellulose filters in the absence of ssDNA.

**Kinetics of ATP Hydrolysis**—Under standard conditions (6 μM recA protein and 50 μM φX ssDNA), recA protein catalyzed ATP hydrolysis with a steady-state rate of 5 min<sup>-1</sup>. Under the same conditions, but with recA1 protein, there was no detectable hydrolysis of ATP (Fig. 4). Although it is not hydrolyzed, ATP does cause the dissociation of recA1 protein from ssDNA. Since ADP also causes the dissociation of recA1 protein from ssDNA, it was conceivable that recA1 protein catalyzed the hydrolysis of a single molecule of ATP to ADP; as a result it dissociated from ssDNA and was somehow inactivated with respect to further ATPase activity. The hydrolysis of a single ATP molecule per recA1 monomer would not be readily detected using the standard TLC-based ATPase assay. Therefore, a more sensitive assay was employed. Reactions were carried out at 50 μM φX ssDNA and 6 μM recA protein using [γ-<sup>32</sup>P]ATP as substrate. At various times, aliquots were quenched with 0.5 N HCl and analyzed for [<sup>32</sup>P]P<sub>i</sub> using the Norit absorption assay.

The pre-steady-state time course for ATP hydrolysis by the wild type recA protein is shown in Fig. 5. Following a lag of approximately 4 s, recA protein catalyzed the hydrolysis of ATP at a rate of 4.8 s<sup>-1</sup> in excellent agreement with the steady-state rate of 5.0 s<sup>-1</sup> obtained using the TLC assay. The lag in the start of ATP hydrolysis was observed under a variety of conditions: (i) when recA protein was added last, (ii) when [γ-<sup>32</sup>P]ATP was added last, and (iii) when [γ-<sup>32</sup>P]ATP was added to an ongoing reaction containing recA protein, ssDNA, and unlabeled ATP. The mechanistic significance of this kinetic lag is under investigation.

The same measurements were made for reactions containing recA1 protein. As shown in Fig. 5, the recA1 protein exhibited no ATPase activity, even when measured at a




FIG. 4. Steady-state ATP hydrolysis by recA and recA1 proteins in the presence of φX ssDNA. Reactions were carried out as described under "Experimental Procedures." Reaction solutions contained 50 μM ssDNA, 200 μM [<sup>3</sup>H]ATP, and 6 μM recA (●) or recA1 protein (○). The points indicated the amount of [<sup>3</sup>H]ADP produced as a function of time as measured by TLC.




FIG. 5. Pre-steady-state ATP hydrolysis by recA and recA1 proteins. Reactions were carried out as described under "Experimental Procedures." Reaction solutions contained 50 μM φX ssDNA and 6 μM recA (●) or recA1 protein (○), with 200 μM [γ-<sup>32</sup>P]ATP added to start the reaction. The points represent the amount of [<sup>32</sup>P]P<sub>i</sub> produced as a function of time as measured using the Norit assay.


sensitivity sufficient to easily detect the hydrolysis of a single equivalent of ATP per recA1 monomer.

The possibility still remained that recA1 protein interacts with ATP to form a covalent phosphoryl-enzyme intermediate that is unable to break down properly, resulting in a modified protein that is inactive with respect to further ATPase activity. If such an intermediate were stable to acid treatment, this event may not have been detected by the Norit assay which detects only <sup>32</sup>P not absorbed to the charcoal. To test for this possibility, ATPase reactions were carried out as described above. After 30 s, the reactions were mixed with trichloroacetic acid to precipitate the protein. The quenched reactions were then filtered on GF/C filters which were then assayed for <sup>32</sup>P. No radioactivity above background (minus protein) was found for either the recA or recA1 proteins (data not shown). Thus, the dissociative effect of ATP on the recA1 protein does not appear to be due to the turnover of a single equivalent of ATP.

**Renaturation of Complementary Single Strands**—The separated (+) and (-) strands of PstI-cleaved [<sup>3</sup>H]φX DNA serve as substrates for the recA protein-promoted renaturation of complementary DNA strands. A typical time course carried out at the optimal ratio (2) of one recA monomer (1 μM) per 30 nucleotides of φX ssDNA (30 μM) is shown in Fig. 6. There was no renaturation during the time course of the experiment in the absence of recA protein. In the presence of recA protein and ATP rapid renaturation occurred with approximately 75% of the ssDNA becoming S1 nuclease-resistant within 10 min. In the absence of ATP, a burst of renaturation was observed (approximately 30%) followed by a slow second phase (*t* > 5 min). When the MgCl<sub>2</sub> concentration was increased from 10 to 30 mM, the ATP-independent reaction proceeded at the same rate as the ATP-stimulated reaction.

The renaturation activity of the recA1 protein was determined under the same conditions used for the wild type protein (Fig. 6). In 10 mM MgCl<sub>2</sub>, the recA1 protein promoted an ATP-independent renaturation reaction identical to that obtained with recA protein. Furthermore, increasing the MgCl<sub>2</sub> concentration from 10 to 30 mM resulted in a stimulation of the ATP-independent renaturation reaction exactly as observed with the wild type protein. However, when the MgCl<sub>2</sub> concentration was kept at 10 mM and ATP (500 μM)

FIG. 6. Renaturation of alkali-denatured [ $^3\text{H}$ ] $\phi\text{X}$  DNA by *recA* and *recA1* proteins. Reactions were carried out as described under "Experimental Procedures." Reaction solutions contained 30  $\mu\text{M}$  denatured [ $^3\text{H}$ ]DNA, 1  $\mu\text{M}$  *recA* protein (left) or 1  $\mu\text{M}$  *recA1* protein (right), and the indicated concentrations of  $\text{MgCl}_2$  and ATP.



was added, the renaturation activity of the *recA1* protein was completely eliminated. This finding is consistent with dissociation of *recA1* protein from ssDNA in the presence of ATP (ADP also eliminated the renaturation activity of *recA1* protein). Thus the *recA1* mutation clearly separates the ATP-dependent and ATP-independent pathways by which the *recA* protein catalyzes the renaturation of complementary DNA strands.

#### DISCUSSION

*RecA1* mutants of *E. coli* are defective in all known functions of the *recA* gene (8). The *recA1* mutation has been identified as a missense mutation in which a glycine at position 160 is replaced by an aspartic acid residue (9). The purified *recA1* protein does not function as a ssDNA-dependent ATPase and is unable to promote DNA strand exchange (3, 10). Nevertheless, it is able to catalyze the ATP-independent renaturation of complementary DNA strands.

The *recA1* protein binds ssDNA with a stoichiometry of 1 monomer/3.5 nucleotides, a value that is very close to 1 monomer/4 nucleotides found for the wild-type protein using nuclease protection (6) as well as other methods (3, 11). The binding of both *recA* and *recA1* proteins to ssDNA appears to be cooperative based on nitrocellulose filter-binding measurements. These results support earlier reports that *recA1* protein forms a complex with ssDNA in the absence of ATP (3, 12).

The affinity of the *recA* and *recA1* proteins for ssDNA were compared by analyzing the kinetics of protein transfer between ssDNA populations. In control reactions with the *recA* protein, transfer was measured by both nuclease protection and nitrocellulose filter binding. Under conditions of excess challenge DNA, the two assays gave similar time courses of equilibration with a  $t_{1/2}$  of about 5 min in the absence of nucleotide. Agreement between the two assays was not necessarily expected. The nuclease protection assay measures total *recA* protein bound to the initial  $^3\text{H}$ -strand; a transfer of 50% of the *recA* monomers results directly in a 50% decrease in nuclease protection. In the nitrocellulose filter-binding assay, however, a transfer of 50% of the *recA* monomers from the initial  $^3\text{H}$ -strand could conceivably leave each  $^3\text{H}$ -labeled DNA molecule with half its *recA* protein; full nitrocellulose filter binding might still occur and the transfer would go

undetected. The finding that the nuclease protection and the filter binding assays *do* give the same time course suggests that transfer of *recA* protein is cooperative with many *recA* monomers (clusters) transferring from one DNA molecule to another in a single event. In this case, at 50% transfer, 50% of the  $^3\text{H}$ -labeled DNA molecules would have lost *all* of their *recA* protein, resulting in a 50% decrease in nuclease protection and a corresponding 50% decrease in filter binding.

Cooperative transfer of the *recA* protein may proceed by the interaction of a *recA*-ssDNA complex with a second ssDNA molecule, resulting in the formation of a transiently two-stranded intermediate. This intermediate could then undergo a "switching" event leading to the transfer of the *recA* cluster from the initial ssDNA molecule to the second ssDNA molecule. Alternatively, transfer may occur by the cooperative dissociation of a cluster of *recA* monomers followed by the cooperative rebinding of the monomers to the second strand.

We were unable to use the nuclease protection assay to measure the transfer of *recA1* protein because of the inability of  $\text{ATP}\gamma\text{S}$  to function as an effective quench for the transfer process.<sup>2</sup> However, transfer of *recA1* protein could be measured by filter binding. The half-time for equilibration using 15 eq of challenge DNA was approximately 1 min, about 5-fold faster than that found for the wild type *recA* protein, indicating that the mutant protein may have a reduced affinity for ssDNA.

Although the *recA* and *recA1* proteins interact similarly with ssDNA in the absence of nucleotide, they exhibit very different properties in the presence of nucleotide cofactors. ATP causes an increase in the rate of transfer of wild type *recA* protein as a result of its hydrolysis to ADP and  $\text{P}_i$  (6). Despite the increase in rate of transfer, nuclease protection experiments indicate that most of the *recA* protein is still bound to ssDNA at any given time; even in the presence of ATP (and an ATP regeneration system), the protection stoichiometry remains at 1 *recA* monomer/4 nucleotides. As noted above, the transfer may occur by a cooperative two-strand switching mechanism that does not necessarily require dissociation of *recA* protein.

We showed previously that  $\text{ATP}\gamma\text{S}$  causes a 2-fold increase in the stoichiometry of nuclease protection by the *recA* protein when ssDNA is in excess relative to *recA* protein. We also

found that addition of ATP $\gamma$ S to ssDNA complexed with subsaturating levels of *recA* protein results in the formation of highly condensed networks of DNA and protein. One interpretation of these results is that ATP $\gamma$ S traps *recA* protein in a conformation in which it is tightly bound to two strands of ssDNA (6). Formation of such networks and the increase in binding stoichiometry necessarily require an excess of ssDNA over *recA* protein. When *recA* protein is in excess over ssDNA, and ATP $\gamma$ S is added, the stoichiometry remains at 1 *recA* monomer/4 nucleotides.<sup>3</sup> In this case there would only be enough ssDNA to occupy a single binding site.

*RecA1* protein also interacts with ATP $\gamma$ S by the following two criteria: (i) direct binding of *recA1*·ssDNA·ATP $\gamma$ S complexes to nitrocellulose filters and (ii) reduction in the rate of transfer of *recA1* protein in the presence of ATP $\gamma$ S. Despite this apparent interaction, ATP $\gamma$ S does not cause the 2-fold increase in the stoichiometry of nuclease protection by the *recA1* protein, nor does it produce the highly condensed *recA1*-ssDNA networks typical of wild type *recA* protein. Thus, the doubled stoichiometry and the network formation found for wild type *recA* protein may be related phenomena.

The pattern of renaturation catalyzed by the *recA* and *recA1* proteins is consistent with these proposals. The wild type *recA* protein catalyzes renaturation in the presence of 10 mM MgCl<sub>2</sub> even in the absence of ATP. We previously suggested that this reaction proceeds via the formation of aggregates of *recA*·ssDNA complexes which serve to bring DNA sequences into a higher effective concentration, thereby increasing the rate of renaturation (2). Such aggregates may form via the interaction of *recA*·ssDNA complexes with a second strand of ssDNA as was suggested above for the *recA* transfer reaction. ATP independent renaturation in the presence of 10 mM Mg<sup>2+</sup> shows a burst of renaturation (approximately 30% of maximum in 1 min) followed by a very slow second phase. The slow phase may be due to the initial formation of a fraction of nonproductive aggregates in which the DNA strands are not in position to renature. ATP may stimulate renaturation by stimulating the movement of *recA* protein to new strands, thereby facilitating the formation of new productive aggregates. Alternatively ATP may stimulate renaturation by facilitating the dissociation of *recA* protein from ssDNA, thereby allowing the zipping of helices to occur unimpeded.

The *recA1* protein shows the same pattern of interaction with ssDNA in the absence of ATP as the wild type protein, and it also shows a similar pattern of ATP-independent renaturation. The *recA1* protein-promoted renaturation reactions are thus also accommodated by a model in which *recA1*·ssDNA complexes interact with a second strand of ssDNA to form renaturation intermediates. Unlike the wild type protein, inclusion of ATP causes dissociation of *recA1*

protein from ssDNA, thereby preventing aggregation and eliminating renaturation activity.

It has recently been reported that the incubation of *recA* protein with ssDNA results in the formation of protein·DNA aggregates that are large enough to be sedimented out of solution by low speed centrifugation (13). These aggregates may correspond to the intermediates in *recA*-promoted renaturation proposed above. These intermediates occur only with subsaturating levels of *recA* protein, consistent with the proposal that they form via ssDNA·*recA*·ssDNA interactions and not ssDNA·*recA*·*recA*·ssDNA interactions. It has also been reported that the *recA1* protein is incapable of forming aggregates with ssDNA (12). This observation was made in the presence of ATP, which we show here causes the dissociation of *recA1* protein from ssDNA and the elimination of renaturation activity. Since *recA1* protein does promote the normal pattern of ATP-independent renaturation, we tested the ability of *recA1* protein to promote the aggregation of ssDNA in the absence of ATP. Our results show that *recA1* protein does indeed cause aggregation of ssDNA in absence of ATP under the same conditions that it promotes renaturation. In agreement with the reported results (12), no aggregation by *recA1* protein was found in the presence of ATP.<sup>2</sup> Thus, the patterns of *recA* and *recA1* protein-induced aggregation of ssDNA are consistent with the patterns of *recA*- and *recA1*-promoted renaturation.

*Acknowledgment*—We gratefully acknowledge a generous gift of *recA1* protein from Dr. Steve West (Yale University).

#### REFERENCES

- Weinstock, G. M., McEntee, K., and Lehman, I. R. (1979) *Proc. Natl. Acad. Sci. U. S. A.* **76**, 126-130
- Bryant, F. R., and Lehman, I. R. (1985) *Proc. Natl. Acad. Sci. U. S. A.* **82**, 297-301
- West, S. C., Cassuto, E., Muraslim, J., and Howard-Flanders, P. (1980) *Proc. Natl. Acad. Sci. U. S. A.* **77**, 2569-2573
- Cox, M. M., McEntee, K., and Lehman, I. R. (1981) *J. Biol. Chem.* **256**, 4676-4678
- Cox, M. M., and Lehman, I. R. (1981) *Proc. Natl. Acad. Sci. U. S. A.* **78**, 3433-3437
- Bryant, F. R., Taylor, A. R., and Lehman, I. R. (1985) *J. Biol. Chem.* **260**, 1196-1202
- Meek, T. D., Johnson, K. A., and Villafranca, J. J. (1982) *Biochemistry* **21**, 2158-2167
- Clark, A. J., and Margulis, A. D. (1965) *Proc. Natl. Acad. Sci. U. S. A.* **53**, 451-459
- Kawashima, H., Horri, T., Ogawa, T., and Ogawa, H. (1984) *Mol. & Gen. Genet.* **193**, 288-292
- Rusche, J. R., Konigsberg, W., and Howard-Flanders, P. (1985) *J. Biol. Chem.* **260**, 949-955
- Morrill, S. W., and Cox, M. M. (1985) *Biochemistry* **24**, 760-767
- Wabiko, H., Ogawa, T., and Ogawa, H. (1983) *Eur. J. Biochem.* **137**, 263-267
- Tsang, S. S., Chow, S. A., and Radding, C. M. (1985) *Biochemistry* **24**, 3226-3232

<sup>3</sup> S. K. Neunendorf and M. M. Cox, personal communication.