

An initial ATP-independent step in the unwinding of a herpes simplex virus type I origin of replication by a complex of the viral origin-binding protein and single-strand DNA-binding protein

Xiaodun He and I. R. Lehman*

Department of Biochemistry, Beckman Center, Stanford University School of Medicine, Stanford, CA 94305-5307

Contributed by I. R. Lehman, January 16, 2001

Using a spectrophotometric assay that measures the hyperchromicity that accompanies the unwinding of a DNA duplex, we have identified an ATP-independent step in the unwinding of a herpes simplex virus type 1 (HSV-1) origin of replication, *Ori_s*, by a complex of the HSV-1 origin binding protein (UL9 protein) and the HSV-1 single-strand DNA binding protein (ICP8). The sequence unwound is the 18-bp A + T-rich segment that links the two high-affinity UL9 protein binding sites, boxes I and II of *Ori_s*. P1 nuclease sensitivity of *Ori_s* and single-strand DNA-dependent ATPase measurements of the UL9 protein indicate that, at 37°C, the A + T-rich segment is sufficiently single stranded to permit the binding of ICP8. Binding of the UL9 protein to boxes I and II then results in the formation of the UL9 protein-ICP8 complex, that can, in the absence of ATP, promote unwinding of the A + T-rich segment. On addition of ATP, the helicase activity of the UL9 protein-ICP8 complex can unwind boxes I and II, permitting access of the replication machinery to the *Ori_s* sequences.

The herpes simplex virus type 1 (HSV-1) origin binding protein (UL9 protein) is a DNA helicase that, when complexed with the HSV-1 single-strand DNA binding protein (ICP8), can unwind an HSV-1 origin (*Ori_s*) to initiate viral DNA replication (1). Measurement of the helicase activity of the UL9 protein has typically made use of nondenaturing polyacrylamide gel electrophoresis (PAGE) to separate the single-stranded DNA products from the duplex DNA substrate (2). This assay, although unambiguous, cannot measure the early events in the unwinding reaction that are necessary to an understanding of its mechanism.

We describe here an optical assay that depends on the hyperchromic effect that results from the rupture of hydrogen bonds as the DNA duplex is unwound. This assay, which can monitor the continuous unwinding of *Ori_s*, has revealed that the initial phase of unwinding of *Ori_s* consists of the UL9 protein and ICP8-dependent, but ATP-independent unwinding of the A + T-rich sequence linking the two inverted repeats, boxes I and II. On addition of ATP, the UL9 protein-ICP8 complex can then promote the bidirectional unwinding of the entire *Ori_s* sequence (3, 4).

Materials and Methods

Nucleotides and Enzymes. [γ -³²P]ATP (6000 Ci/mmol) and proteinase K were obtained from DuPont/NEN and Roche Molecular Biochemicals, respectively. *Escherichia coli* single-stranded DNA binding protein (SSB), P1 nuclease, and T4 polynucleotide kinase were from United States Biochemical. Recombinant UL9 protein and ICP8 were purified as previously described (3, 5). They were free of detectable exo- and endonuclease activity.

Oligonucleotides. Oligonucleotides were purchased from Operon Technologies (Alameda, CA). They were purified by 16% PAGE

under denaturing conditions. Poly(dT)_{300–3000} was obtained from Sigma. When required, the oligonucleotides were labeled with [γ -³²P]ATP with polynucleotide kinase according to a protocol supplied by Amersham Pharmacia. Unincorporated nucleotide was removed by G-50 spin column chromatography (Quickspin DNA; Roche Molecular Biochemicals).

***Ori_s*.** The complementary oligonucleotides: 5'-CGCGAAGCGTTCGCACTTCGTCCCAATATATATATATTATTAGGGCGCAAGTGCAGCACTGGC-3', and 5'-GCCAGTGCTCGCACTTCGCCCTAATAATATATATATATTGGGACGAA-GTGCGAACGCTTCGCG-3', both at 216 μ g/ml in 50 mM Tris-HCl (pH 8.0) containing 0.1 M NaCl were mixed, heated at 90°C for 5 min, then slowly cooled to room temperature. The *Ori_s* duplex that was generated lacked Box III.

***Ori_sT₁₈*.** The complementary oligonucleotides: 5'-CGCGAAGCGTTCGCACTTCGTCCCTTTTTTTTTTTTTTTTTTTT-TGGGCGAAGTGCAGCACTGGC-3', and 5'-GCCAGTGCTCGCACTTCGCCCTTTTTTTTTTTTTTTTTTTTGGGACGAAGTGCAGAACGCTTCGCG-3', both at 197 μ g/ml, were annealed as described above for *Ori_s* to generate an *Ori_s* analogue in which the A + T-rich segment was replaced by a "bubble" consisting of two 18-nt oligodeoxythymidylates.

***Ori_s(GC)₁₈*.** The complementary oligonucleotides: 5'-CGCGAAGCGTTCGCACTTCGTCCCGGCGCGCGCGCGCGCGCGCGCGCGGAAGTGCAGCACTGGC-3', and 5'-GCCAGTGCTCGCACTTCGCCCGGCGGGCGCGCGCGCGCGCGCGGACGAAGTGCAGAACGCTTCGCG-3', both at 197 μ g/ml, were annealed as described above for *Ori_s*, to generate an *Ori_s* analogue in which the A + T = rich segment was replaced by (GC)₁₈.

Mutant *Ori_s*. The complementary oligonucleotides: 5'-GCGAAGCGGGCGCACTTCGTCCCAATATATATATATATTATTAGGGCGAAGTGCAGCACTGGC-3', and 5'-GCCAGTGCTCGCACTTCGCCCTAATAATATATATATATTGGGACGAAAGTGCAGAACGCTTCGCG-3', both at 144 μ g/ml, were annealed as described above for *Ori_s*.

Box I. The complementary oligonucleotides: 5'-CGCGAAGCGTTCGCACTTCGTCCC-3', and 5'-CGGACGAAGTGCAGAACGCTTCGCG-3', both at 41 μ g/ml, were annealed as described for *Ori_s*.

Abbreviations: HSV-1, herpes simplex virus type 1; SSB, single-stranded DNA binding protein; *T_m*, melting temperature.

*To whom reprint requests should be addressed. E-mail: blehman@cmgm.stanford.edu.
The publication costs of this article were defrayed in part by page charge payment. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. §1734 solely to indicate this fact.

Spectrophotometric Assay of Helicase Activity. Absorbance at 260 nm was measured with a Uvicom (Kontron, Zurich) UV/VIS spectrophotometer using a quartz cuvette with a path length of 1 cm. The cuvette compartment was kept at 37°C, and absorbance measurements were made at 0.5-s intervals for 30 min. Initial velocities ($\Delta A_{260}/\text{min}$) were derived from the linear part of the curve. The reaction mixture (1 ml) contained 50 mM Tris-HCl (pH 8.15), 10 mM NaCl, 10 mM MgCl₂, 1 mM DTT, 0.01% Nonidet P-40, Ori_s, mutant Ori_s, Ori_sT₁₈, or Ori_s(GC)₁₈, UL9 protein, and/or ICP8 as indicated. The reference cuvette contained the same reaction mixture without UL9 protein or ICP8.

Measurement of P1 Nuclease Sensitivity. The reaction mixtures (25 μl) contained 50 mM Tris-HCl (pH 8.15), 10 mM NaCl, 10 mM MgCl₂, 2 mM DTT, 0.01% Nonidet P-40, and 1 pmol of ³²P-labeled Ori_s or Ori_s T₁₈. Increasing amounts of P1 nuclease (5×10^{-6} units/ μl) containing 2.5 mM ZnCl₂ were added, and the mixtures were incubated at either 25°C or 37°C for 5 min.

The reactions were terminated by the addition of 2 μl of 0.5 M EDTA. An aliquot (8 μl) was loaded together with 80% formamide loading buffer containing 0.05% bromophenol blue and 0.05% xylene cyanol onto a 300 \times 400 \times 0.4-mm 10% polyacrylamide sequencing gel containing 8 M urea and 40% formamide that had been prerun until the temperature of the gel was 50°C (1–2 h). After electrophoresis, the gel was dried on a Whatman No. 1 filter. The relative intensity of the bands was measured with a PhosphorImager (Molecular Dynamics).

DNA-Dependent ATPase Assay. Reaction mixtures (25 μl) contained Tris-HCl (pH 8.15), 10 mM MgCl₂, 1 pmol UL9 protein, 1 pmol ICP8, 10 μg BSA, poly(dT)_{300–3000}, Box I, Ori_s, Ori_sT₁₈, Ori_s(GC)₁₈, or mutant Ori_s, 1 μC of [γ -³²P]ATP, and 3 mM ATP. Mixtures were incubated at 37°C for 1 h. Reactions were stopped by the addition of 3 μl of 0.5 M EDTA, pH 8.0, and chilled on ice. Aliquots (1 μl) were spotted onto polyethyleneimine-cellulose F, TLC plates and dried. The ATP substrate and inorganic phosphate product were separated with 1.0 M formic acid/0.4 M LiCl as solvent. The TLC plates were quantitated with a PhosphorImager (Molecular Dynamics).

DNA Helicase Assay. Reaction mixtures (25 μl) containing 50 mM Hepes-KOH (pH 8.15), 10 mM NaCl, 10 mM MgCl₂, 2.0 mM DTT, 10% (vol/vol) glycerol, 10 μg BSA, 1.0 pmol Ori_sT₁₈, or 1.0 pmol Ori_s(GC)₁₈, and the indicated amounts of ICP8 and UL9 protein were incubated on ice for 5 min. ATP (50 μM) was added to the reaction mixtures, which were then incubated at 37°C for 60 min. The reactions were terminated by the addition of 6.5 μl of stop solution (100 mM EDTA/1% SDS/20 μg proteinase K), then subjected to electrophoresis through a 15% polyacrylamide gel at 10 V/cm. The gel was dried on Whatman DE81 paper at 80°C under vacuum and quantitated with a PhosphorImager (Molecular Dynamics) (4).

Gel Mobility Shift Analysis. Reaction mixtures (50 μl) containing 50 mM Hepes-KOH (pH 8.15), 2.0 mM DTT, 50 mM NaCl, 10 mM MgCl₂, 10 μg BSA, 10% (vol/vol) glycerol, and the indicated amounts of UL9 protein and ICP8 were preincubated on ice for 30 min. 5'-³²P-labeled Ori_s(GC)₁₈ (0.5 pmol) was added, and the mixtures were incubated at 37°C for 15 min. Four percent nondenaturing PAGE of the reaction mixtures was performed at room temperature with Tris-HCl/glycine buffer containing 50 mM Tris-HCl and 380 mM glycine buffer at 30 V and 25 mA for 2.5 h. After electrophoresis, the gel was dried on Whatman DE81 paper at 80°C under vacuum and visualized by a PhosphorImager (Molecular Dynamics) (6).




Fig. 1. Spectrophotometric measurement of unwinding of Ori_s. Reaction mixtures prepared as described in *Materials and Methods* contained 50 nM Ori_s(1), or mutant Ori_s(2), 4 nM UL9 protein, and 5 nM ICP8. Increase in A_{260} was measured as described in *Materials and Methods*.

Results

Spectrophotometric Assay for Unwinding of Ori_s. The increase in A_{260} as a consequence of the unwinding of Ori_s by the UL9 protein-ICP8 complex is shown in Fig. 1. The reaction proceeded linearly for 40 min. ATP was not required for the reaction, and there was no effect of adding the nonhydrolyzable ATP analogue ATP γ S (data not shown). The initial rate was linearly dependent on the concentration of UL9 protein and ICP8 (Figs. 2 and 3). However, the same plateau value was reached over a 5-fold range of UL9 protein-ICP8 concentrations (4–20 pmol; data not shown). ICP8 could not be replaced by the heterologous *E. coli* single-strand DNA binding protein, SSB (Fig. 3). A mutant Ori_s with mutations in Boxes I and II that was unable to bind UL9 protein (7) showed no increase in A_{260} (Fig. 1). The reaction depended on Mg²⁺ and was optimal at 10–20 mM Mg²⁺. In agreement with earlier measurements of helicase activity by nondenaturing PAGE, the reaction was optimal at 40–45°C and declined rapidly at higher temperatures, presum-




Fig. 2. Effect of UL9 protein concentration on initial rate of unwinding of Ori_s. Reaction mixtures were prepared as described in *Materials and Methods* with 320 nM Ori_s, 13.2 nM ICP8, and the indicated concentrations of UL9 protein. Initial rate units ($\Delta A_{260}/\text{min}$) were determined from the increase in A_{260} at 15 min.




Fig. 3. Effect of ICP8 concentration on initial rate of unwinding of *Ori_s*. Reaction mixtures were prepared as described in *Materials and Methods* with 320 nM *Ori_s*, 5 nM UL9 protein, and the indicated concentrations of ICP8 (\blacktriangle) or *E. coli* SSB (\blacksquare).

ably as a result of thermal denaturation of the enzyme (2) (Fig. 4). The T_m for the reaction was 37°C. The inability to observe any reaction at 30°C suggests that some thermally induced destabilization of the A + T-rich sequence is necessary for the reaction (see below).

Unwinding of A + T-Rich Sequence Is the Initial Step in Unwinding of *Ori_s*. The thermal denaturation profile of *Ori_s* is shown in Fig. 5. The T_m of 80°C is consistent with the G + C content of *Ori_s* (49%). The maximum increase in A_{260} that occurred on incubation of *Ori_s* with the UL9 protein and ICP8 was 23% of that obtained on complete thermal denaturation (Figs. 1 and 5). Thus, as measured by the spectrophotometric assay, 23% of the *Ori_s* duplex was unwound by the UL9 protein-ICP8 complex. This value is in good agreement with the proportion of *Ori_s* comprising the A + T-rich sequence (18/63 = 28%), suggesting that it is this sequence that is being unwound in the initial




Fig. 4. Effect of temperature on initial rate of unwinding of *Ori_s*. Reaction mixtures were prepared as described in *Materials and Methods* with 52 nM *Ori_s*, 5.6 nM UL9 protein, and 5.2 nM ICP8. Temperature was varied from 10°C to 55°C as indicated.




Fig. 5. Temperature melting curve of *Ori_s*. DNA melting was measured with a Cary IE UV-visible spectrophotometer equipped with a temperature control unit. The temperature was increased or decreased at a rate of 1°C per min and held at that temperature for 30 min. The reaction cuvette contained 500 nM *Ori_s* in 50 mM Tris-HCl (pH 8.15). The reference cuvette contained the Tris buffer without *Ori_s*. The temperature ranges were 20° to 90°C (\blacktriangle) and 90° to 20°C (\square).

ATP-independent phase of the reaction. This idea was supported by an experiment using an *Ori_s* analogue in which the A + T-rich sequence was replaced by two 18-nt oligodeoxythymidylate sequences to create an unpaired “bubble” between Boxes I and II (*Ori_sT₁₈*) (4). As shown in Fig. 6, there was no increase in A_{260} with *Ori_sT₁₈* under conditions in which an increase of 23% of maximum was observed with *Ori_s*. Because the oligodeoxythymidylate sequences in *Ori_sT₁₈* were already unpaired, no hyperchromic effect was to be expected.

Further evidence that a thermally induced destabilization of the A + T-rich sequence is required for the initial ATP-independent step in unwinding of *Ori_s* was provided by an experiment in which the A and T residues of the A + T-rich sequence were replaced by G and C [*Ori_s(GC)₁₈*]. With this *Ori_s* analogue in which the G + C sequence should remain fully hydrogen bonded at 37°C, there was no increase in A_{260} (Fig. 6).




Fig. 6. Lack of hyperchromic effect with *Ori_sT₁₈* and *Ori_s(GC)₁₈*. Reaction mixtures were prepared as described in *Materials and Methods* with 340 nM *Ori_sT₁₈* and 4 nM UL9 protein (1), 340 nM *Ori_sT₁₈* and 5 nM ICP8 (2), 340 nM *Ori_sT₁₈* and 4 nM UL9 protein, and 5 nM ICP8 (3) or 53 nM *Ori_s(GC)₁₈*, 4 nM UL9 protein, and 5 nM ICP8 (4).




Fig. 7. Lack of ATP-dependent unwinding of $\text{Ori}_S(\text{GC})_{18}$ by UL9 protein-ICP8 complex. (a) Reaction mixtures for helicase measurements were prepared as described in *Materials and Methods* with 1 pmol $\text{Ori}_S\text{T}_{18}$ or 1 pmol $\text{Ori}_S(\text{GC})_{18}$, and the indicated amounts of UL9 protein and ICP8. Δ , $\text{Ori}_S\text{T}_{18}$ and $\text{Ori}_S(\text{GC})_{18}$, heated at 100°C for 5 min. Incubation and PAGE were performed as described in *Materials and Methods*. (b) Reaction mixtures for gel mobility shift measurements were prepared as described in *Materials and Methods* with 0.5 pmol $\text{Ori}_S(\text{GC})_{18}$ and the indicated amounts of UL9 protein and ICP8. Samples were processed as described in *Materials and Methods*. ds, double stranded; ss, single stranded.

Furthermore, unlike $\text{Ori}_S\text{T}_{18}$, which could be efficiently unwound by the helicase activity of the UL9 protein-ICP8 complex (4), $\text{Ori}_S(\text{GC})_{18}$ was essentially inert (Fig. 7). The lack of helicase activity was not a consequence of the failure of UL9 protein to bind $\text{Ori}_S(\text{GC})_{18}$ as judged by a gel mobility shift analysis (Fig. 7).

Single-Stranded Character of A + T-Rich Sequence. P1 nuclease has a strong preference for single- over double-stranded DNA (8).




Fig. 9. DNA-dependent ATPase activity of UL9 protein. Reactions were performed as described in *Materials and Methods* with 1 pmol UL9 protein, 1 pmol ICP8, 3 mM $[\gamma\text{-}^{32}\text{P}]\text{ATP}$, and 7.4 pmol poly(dT), 5.7 pmol Box I, 13 pmol Ori_S , 11 pmol $\text{Ori}_S\text{T}_{18}$, 7.7 pmol mutant Ori_S or $\text{Ori}_S(\text{GC})_{18}$.

Treatment of $\text{Ori}_S\text{T}_{18}$ with P1 nuclease at both 25°C and 37°C produced a digestion pattern consistent with the presence of oligo(dT)₁₈. In contrast, the pattern observed on nuclease digestion of the wild-type Ori_S duplex at 37°C indicated that the A + T sequence was largely destabilized at 37°C but not at 25°C. As expected, Boxes I and II of neither substrate were P1 nuclease-sensitive at either temperature (Fig. 8).

Further support for the single-stranded character of the A + T-rich sequence was provided by measurement of the single-




Fig. 8. P1 nuclease sensitivity of Ori_S and $\text{Ori}_S\text{T}_{18}$. Reaction mixtures prepared as described in *Materials and Methods* with 1 pmol of ^{32}P -labeled Ori_S or $\text{Ori}_S\text{T}_{18}$ and the indicated amounts of P1 nuclease and incubated at 25°C or 37°C. The products were analyzed by 10% PAGE in the presence of urea-formamide as described in *Materials and Methods*. M, markers.

strand DNA-dependent ATPase activity of the UL9 protein (9). As shown in Fig. 9, although the fully duplex Box I supported little ATPase activity at 37°C, Ori_s showed substantial ATPase activity, similar to that observed with Ori_sT₁₈ and with poly(dT)_{300–3000}. In contrast, only a low level of DNA-dependent ATPase activity was observed with Ori_s at 25°C, presumably a manifestation of the DNA-independent ATPase activity of the UL9 protein (9) (data not shown). A mutant Ori_s that is unable to bind the UL9 protein did not support the ATPase activity of the UL9 protein (Fig. 9). Similarly, no DNA-dependent ATPase activity was observed with Ori_s(GC)₁₈.

Discussion

The spectrophotometric assay that we have developed which permits the continuous monitoring of helicase activity has provided some insight into the initial steps in the unwinding of Ori_s by the UL9 protein–ICP8 complex. The A + T-rich sequence linking boxes I and II of Ori_s, because of its low *T_m* relative to the G + C-rich boxes I and II, possesses sufficient single-strand character at 37°C to permit the binding of ICP8. However, complete destabilization of the duplex A + T-rich sequence does not occur under these conditions, as judged by the lack of hyperchromicity on ICP8 binding. The UL9 protein can, however, bind to the high-affinity boxes I and II and form a complex with ICP8 bound to the A + T-rich sequence. The resulting UL9 protein–ICP8 complex, presumably using the energy of binding, can then completely unwind the A + T-rich sequence in an ATP-independent reaction. As observed previ-

ously, a specific UL9 protein–ICP8 complex is required (2–4); the *E. coli* SSB, although capable of binding to the A + T-rich sequence (10), is unable to promote its unwinding, presumably because of its inability to interact with the UL9 protein. In the presence of an energy source, i.e., ATP, the UL9 protein–ICP8 complex can promote the complete unwinding of Ori_s to permit access of the replication machinery to the origin sequences.

We had earlier suggested that the unwinding of the A + T-rich sequence depended on transcription from the transactivator binding sequences flanking Ori_s, i.e., transcriptional activation (3). Although transcription may indeed activate the initiation of HSV-1 DNA replication at Ori_s *in vivo*, the findings reported here suggest that this is not an obligatory event.

Earlier experiments by Koff *et al.* (11) had shown that the A + T-rich sequence at the HSV-2 Ori_s became DNase I sensitive on binding of the UL9 protein. In KMnO₄-sensitivity measurements, they also observed the increased reactivity of the thymidylate residues of the A + T sequence on addition of the UL9 protein. Based on these findings, they suggested that binding of UL9 to Boxes I and II resulted in the destabilization of the A + T sequence. Our studies have indicated that the A + T sequence is partially unwound at 37°C, even in the absence of UL9 protein. This reaction might, however, be facilitated by the binding of UL9 protein.

We thank Drs. Steven Matson, Per Elias, and Paul Boehmer for their comments and suggestions. This work was supported by National Institutes of Health Research Grant AI26538.

1. Lehman, I. R. & Boehmer, P. E. (1999) *J. Biol. Chem.* **274**, 28059–28062.
2. Boehmer, P. E., Dodson, M. S. & Lehman, I. R. (1993) *J. Biol. Chem.* **268**, 1220–1225.
3. Lee, S. S.-K. & Lehman, I. R. (1997) *Proc. Natl. Acad. Sci. USA* **94**, 2838–2842.
4. He, X. & Lehman, I. R. (2000) *J. Virol.* **74**, 5726–5728.
5. Boehmer, P. E. & Lehman, I. R. (1993) *J. Virol.* **67**, 711–715.
6. Lee, S. S.-K., Dong, Q., Wang, T. S.-F. & Lehman, I. R. (1995) *Proc. Natl. Acad. Sci. USA* **92**, 7882–7886.
7. Hernandez, T. R., Dutch, R. E., Lehman, I. R., Gustafsson, C. & Elias, P. (1991) *J. Virol.* **65**, 1649–1652.
8. Chiu, S. K., Rao, B. J., Staory, R. M. & Radding, C. M. (1993) *Biochemistry* **32**, 13246–13155.
9. Dodson, M. S. & Lehman, I. R. (1993) *J. Biol. Chem.* **268**, 1213–1219.
10. Makhov, A. M., Boehmer, P. E., Lehman, I. R. & Griffith, J. D. (1996) *EMBO J.* **15**, 1742–1750.
11. Koff, A., Schwedes, J. F. & Tegtmeyer, P. (1991) *J. Virol.* **65**, 3284–3292.