

## Energy-Dependent Activation of the Temperature-Sensitive DNA Polymerases Induced by Bacteriophage T4 Gene 43 Mutants<sup>1</sup>

JEREMY THORNER,<sup>2</sup> WAI MUN HUANG,<sup>3</sup> AND I. R. LEHMAN

*Department of Biochemistry, Stanford University School of Medicine, Stanford, California 94305*

*Accepted June 27, 1975*

Temperature shift experiments with temperature-sensitive mutants in T4 gene 43 indicate that the production of a functional T4 DNA polymerase involves the energy-dependent conversion of an inactive precursor to the active enzyme. Comparisons of the inactive precursor with the active polymerase have demonstrated that the energy-dependent activation affects both the polymerase and 3' → 5' exonuclease activities of the enzyme. Despite evidence suggesting that the energy source for the activation process is a high-energy phosphate compound, the energy-dependent step does not appear to involve covalent modification of the enzyme.

The DNA polymerase synthesized after infection of *Escherichia coli* with bacteriophage T4 is absolutely essential for the initiation and maintenance of viral DNA replication (Epstein *et al.*, 1963; Warner and Hobbs, 1967). As an approach toward understanding more precisely the role of T4 DNA polymerase in the replication of T4 DNA, we examined the DNA polymerase induced after infection of *E. coli* with the T4 mutant, *ts* L56, which bears a temperature-sensitive mutation in gene 43, the structural gene for T4 DNA polymerase (de Waard *et al.*, 1965). In the course of these studies, a novel phenomenon was encountered. It was found that at the nonpermissive temperature, an inactive form of the polymerase was synthesized, which could be converted to the functional enzyme upon lowering the temperature. The ap-

pearance of active enzyme after temperature stepdown was not prevented by inhibitors of protein synthesis, but was blocked by anaerobiosis or by inhibitors of energy-yielding metabolism (Swartz *et al.*, 1972). The fact that the appearance of active enzyme was energy-dependent clearly distinguished the effect from facile renaturation of a thermolabile protein. Thus, the behavior of the *ts* L56 mutant suggested the involvement of an activation process in the formation of functional T4 DNA polymerase that had previously gone undetected.

In this report, we present experiments that support and extend our initial observations and that attempt to define the molecular basis of the activation process.

### MATERIALS AND METHODS

*Chemicals and enzymes.* Reagent chemicals were obtained from the following sources: ribo- and deoxyribonucleoside triphosphates from Pabst Laboratories; triethanolamine from Fisher; Tris<sup>4</sup> from Sigma; chloramphenicol from Parke-Da-

<sup>1</sup> This work was supported in part by a grant from the United States Public Health Service (GM 06196). We are very grateful to Janice Chien for her help in performing several of the experiments.

<sup>2</sup> Aided by a fellowship from the Jane Coffin Childs Memorial Fund for Medical Research. Present address: Department of Bacteriology and Immunology, University of California, Berkeley, California 94720.

<sup>3</sup> Present address: Department of Microbiology, University of Utah School of Medicine, Salt Lake City, Utah.

<sup>4</sup> Abbreviations: Tris, tris (hydroxymethyl) aminomethane; Brij 58, polyoxyethylene-20-cetyl ether; BSA, bovine serum albumin; dTTP, deoxythymidine 5'-triphosphate; and m.o.i., multiplicity of infection.

vis; Brij 58 from Atlas Chemicals; salt-free lysozyme from Worthington; BSA (fatty acid-free) from Pentex; salmon sperm DNA from Calbiochem; sodium dodecyl sulfate from BDH Chemicals; and media constituents from Difco. Radiochemicals were purchased from the following suppliers: [ $^3\text{H}$ ]dTTP from Schwarz/Mann; [ $^{35}\text{S}$ ]H $_2$ SO $_4$ , [ $^{32}\text{P}$ ]H $_3$ PO $_4$ , and [ $^{125}\text{I}$ ]NaI from New England Nuclear. All other chemicals were reagent grade. Alkali-denatured salmon sperm DNA and nicked heat-denatured [ $^3\text{H}$ ]DNA from *E. coli* were prepared as previously described (Huang and Lehman, 1972a). Antiserum directed against T4 DNA polymerase was kindly prepared for us by Dr. L. Levine. Antiserum directed against *E. coli* exonuclease I was obtained as described by Aposhian and Kornberg (1962). Antisera were diluted into 0.15 M NaCl and heated at 70° for 30 min prior to use in order to inactivate nuclease activity. Wild-type T4 DNA polymerase and the *ts* L56 mutant enzyme were purified as previously described (Hall and Lehman, 1968).

*Bacterial and bacteriophage strains and growth conditions.* *E. coli* strains B (wild-type) and CR63 (*sup* D $^+$ )<sup>5</sup> were originally obtained from Drs. R. S. Edgar and J. Drake. Strain JT1200 is a *pol*A $^-$  derivative of *E. coli* B constructed in the following way. A spontaneous *spc* $^r$  clone of *E. coli* B was lightly mutagenized with ethylmethane sulfonate as described by Miller (1972), and a *rha* $^-$  derivative was picked from a rhamnose-tetrazolium indicator plate. A spontaneous methyl-methane sulfonate-resistant (MMS $^r$ ) clone of the mutagenized strain was then selected and presumably represents a revertant to *lon* $^+$ . (The uv- and MMS-sensitivity of wild-type *E. coli* B are manifestations of its *lon* $^-$  character.)<sup>6</sup> This MMS $^r$  derivative was then mated with *E. coli* K12 strain JG78 (Hfr KL-25, *met* E $^-$ , *pol* A1, *rha* $^+$ , *spc* $^s$ ) obtained from Dr. I. Herskowitz. The *rha* $^+$  recombinants were selected on rhamnose-minimal salt plates containing spectino-

mycin and methionine, and then were scored for their methionine requirement, and for the presence of the *pol* A1 mutation by their uv- and MMS-sensitivity. Finally, sonic extracts were prepared from the recombinants with the expected phenotype and were assayed for DNA polymerase I activity as previously described (Lehman and Chien, 1972). JT1200 (*met* E $^+$ , *pol* A1, *rha* $^+$ , *spc* $^r$ ) had less than 0.3% of the DNA polymerase I activity of either its immediate parent or *E. coli* B. It grew in broth with essentially the same doubling time and to the same cell density as either its immediate parent or *E. coli* B. *E. coli* K12 strain AN120 (*unc* A $^-$ ) and its isogenic parent AN180 have been described (Butlin *et al.*, 1971) and were obtained from Dr. F. Gibson.

Temperature-sensitive mutants of bacteriophage T4: L107, L97, L56, S9, L88, L141, and L91 (all in gene 43); and *amber* mutants: N82 (gene 44) and E315 (gene 32), were originally obtained from Drs. R. S. Edgar and J. Drake. Double mutants were constructed by standard crossing procedures (Hayes, 1968).

Media (H-broth and EHA-agar plates) used for the growth of cells and for the propagation and assay of T4 phages were those described by Steinberg and Edgar (1967). In some experiments, HT-broth or HP-broth (H-broth buffered at pH 7.0 with 50 mM triethanolamine hydrochloride or 50 mM potassium phosphate, respectively) were used. Media were supplemented with 0.2 mM arginine and 0.2  $\mu\text{M}$  thiamine for experiments with *E. coli* K-12 strains AN120 and AN180.

*Preparation of cell extracts and enzyme assays.* Cultures (150–300 ml) were grown in 1-liter flasks with vigorous aeration in a gyratory water bath at the appropriate temperature. At various times after phage infection, portions (25 ml) of the infected culture containing about  $5 \times 10^8$  cells/ml were removed from the growth flask as rapidly as possible and immediately added to an approximately equal volume of chipped ice containing 1.5 ml of 1.0 M NaN $_3$ . These precautions prevented spontaneous activation of T4 DNA polymerase during preparation of the extract. After all

<sup>5</sup> Genetic nomenclature is according to Taylor and Trotter (1972).

<sup>6</sup> D. Freifelder, personal communication.

samples were collected, they were centrifuged at 8000 g for 10 min and the cell pellets were resuspended in 2.0 ml of 50 mM Tris-HCl, pH 7.5, containing 10 mM 2-mercaptoethanol. The cells in suspension were then ruptured at 0° by two 30-sec bursts of sonic irradiation from a Mullard sonicator. After the extracts were clarified by centrifugation for 20 min at 12,000 g, the supernatant fluid was removed and was assayed for T4 DNA polymerase activity at 30° essentially as described by DeWaard *et al.* (1965) or by the method of Brutlag and Kornberg (1972). Under either assay condition, activity due to *E. coli* DNA polymerases II and III is negligible (Kornberg and Geftter, 1971, 1972). The 3' → 5' exonuclease activity of T4 DNA polymerase was assayed as reported previously (Huang and Lehman, 1972a). Brij lysates were prepared with only minor modifications of the procedure described before (Swartz *et al.*, 1972).

*Other methods.* Protein concentration was determined by the technique of Lowry *et al.* (1951), using BSA as a standard. Preparation of <sup>35</sup>S-labeled phage-infected cells, electrophoresis in slabs of polyacrylamide containing sodium dodecyl sulfate, and radioimmune assay of T4 DNA polymerase protein were all performed essentially as described previously (Huang and Lehman, 1972b). Microdetermination of phosphate was performed by minor modification of the method of Chen *et al.* (1956).

## RESULTS

*Increase in DNA polymerase activity after downshift is due to the phage-induced enzyme.* It was possible that in our earlier experiments (Swartz *et al.*, 1972) the presence of host DNA polymerase I obscured events related to the production of T4 DNA polymerase after infection. To avoid this problem, a *pol A1* derivative of *E. coli* B was constructed. Infection of this strain with T4 *ts* L56 confirmed our earlier observations. As shown in Fig. 1, only a small amount of T4 DNA polymerase activity appeared after infection at 43°. However, upon lowering the temperature to 30°, a marked and sustained increase in T4 DNA polymerase activity occurred almost immediately. This increase cannot be due to




Fig. 1. Time course of T4 *ts* L56 DNA polymerase appearance in a *polA1* mutant of *E. coli* B. A 200-ml culture of JT1200 growing at 43° in HT-broth was infected at an  $A_{595}$  of 0.8 with T4 *ts* L56 at an m.o.i. of 5. At 25 min after infection, chloramphenicol (CAM) was added to a final concentration of 150  $\mu$ g/ml. Five minutes later the culture was shifted to 30°. Samples were removed at the times indicated, and extracts were prepared and assayed for T4 DNA polymerase activity and for protein as described under Materials and Methods. Extracts were assayed in the absence (O—O) and in the presence (●—●) of 30  $\mu$ l of diluted antiserum directed against purified T4 DNA polymerase.

the synthesis of new T4 DNA polymerase molecules since it was shown earlier that such synthesis is completely prevented by the addition of chloramphenicol (Swartz *et al.*, 1972). That the increase in DNA polymerase activity is indeed due to the T4-induced enzyme was demonstrated by the fact that there was essentially no activity if the extracts were assayed in the presence of antiserum directed against purified T4 DNA polymerase.

*Appearance of T4 DNA polymerase activity after downshift requires unperturbed energy-yielding metabolism.* What distinguishes the production of active enzyme after temperature stepdown from a simple rapid renaturation of the thermolabile protein is its sensitivity to 2,4-dinitrophenol,

an uncoupler of oxidative phosphorylation. Half-maximal inhibition of the appearance of T4 DNA polymerase activity occurred at  $55 \mu\text{M}$  and essentially complete inhibition was achieved at  $0.3 \text{ mM}$  when *E. coli* B was used as the host (Fig. 2). About tenfold higher concentrations of the uncoupler were needed to prevent completely the appearance of active enzyme when JT1200 or an *E. coli* K12 strain was used, probably because of the lower permeability to the drug in the latter strains. The appearance of active enzyme could also be blocked by shifting the infected cells from 43 to  $4^\circ$ . The cells could be maintained at this low temperature for considerable periods of time, with active enzyme appearing only when the culture was returned to  $30^\circ$  (Fig. 3). Earlier studies showed that anaerobiosis or inhibitors of cytochrome oxidase ( $\text{CN}^-$  and  $\text{N}_3^-$ ) also prevented appearance of active enzyme (Swartz *et al.*, 1972). The appearance of functional T4 DNA polymerase activity after temperature-downshift has been termed "activation."

*Activation of T4 DNA polymerase activity is not an allele-specific effect.* Six other




FIG. 2. Effect of various concentrations of 2,4-dinitrophenol on the time course of T4 DNA polymerase formation. A 450-ml culture of *E. coli* B growing at  $43^\circ$  in H-broth was infected at an  $A_{595}$  of 0.8 with T4 *tsL56* at an m.o.i. of 5. The experiment was performed as described for Fig. 1, except that at the time of temperature shift the culture was divided into 50-ml portions to which 2,4-dinitrophenol was added to the final concentrations indicated.




FIG. 3. Effect of low temperature on the time course of T4 DNA polymerase appearance. A 225-ml culture of *E. coli* B growing at  $43^\circ$  in HT-broth was infected at an  $A_{595}$  of 0.8 with T4 *tsL56* at an m.o.i. of 5. The experiment was performed as described for Fig. 1, except that at the time of temperature shift a 50-ml sample of the culture was placed at  $30^\circ$  while the remainder of the culture was placed at  $5^\circ$ . Thirty-five minutes after the first temperature shift, another 50-ml sample of the culture was withdrawn from the  $5^\circ$  culture and placed at  $30^\circ$ , while the remainder was left at the  $5^\circ$  temperature.

temperature-sensitive mutations of T4 DNA polymerase that are located throughout gene 43 (Fig. 4) were tested for activation. As shown in Fig. 5, mutants L107, L97, and S9 behaved like L56: T4 DNA polymerase activity increased ten- to thirty-fold after temperature stepdown. Furthermore, the increase in activity for each of these mutants was completely prevented by 2,4-dinitrophenol. Mutants *tsL88* showed a slight increase in T4 DNA polymerase activity over that present immediately before downshift, which was blocked by 2,4-dinitrophenol. Mutants *tsL91* and *tsL141* showed no significant increase in T4 DNA polymerase activity upon temperature shiftdown.

*Activation does not require a functional "replication complex."* Purified T4 DNA polymerase has been shown to interact *in vitro* with purified preparations of the single-stranded DNA binding protein that is the product of T4 gene 32 (Huberman *et al.*, 1971). Moreover, recent experiments indicate that the absence of T4 DNA polymerase prevents the binding of several other T4 proteins to DNA-cellulose columns, suggesting that the interaction with




FIG. 4. Relationship between the genetic and physical maps of T4 gene 43. Order of the *amber* fragments with respect to the complete protein are according to Huang and Lehman (1972), based on a native molecular weight of 102,000 (Nossal and Hershfield, 1971). Order of the temperature-sensitive mutants with respect to the *amber* mutants is derived from the data of Allen *et al.* (1970). Relative activation refers to T4 DNA polymerase activity at 65 min after temperature shiftdown divided by the T4 polymerase activity just before temperature shift.

the polymerase is necessary for their binding to DNA (Huang and Buchanan, 1974). It also has been demonstrated both *in vivo* and *in vitro* that at least six T4 gene products are required for T4 DNA replication and two of these proteins, the products of genes 44 and 62, have been purified and shown to exist as a complex (Epstein *et al.*, 1963; Barry and Alberts, 1972). However, with double mutants *ts* L56 *am* N81 and *ts* L56 *am* E315 in which DNA synthesis was completely blocked by *amber* mutations in gene 44 or in gene 32, respectively, the formation of active T4 DNA polymerase upon temperature stepdown occurred identically to that found for *ts* L56 alone. These observations suggest that activation can occur even when the polymerase is not part of a functional replication complex.

*An inactive precursor is synthesized at the nonpermissive temperature.* The finding that the increase in T4 DNA polymerase activity that occurs upon temperature shift down does not depend on protein synthesis suggests that the molecules that are activated subsequently to form the functional enzyme have already been made at the restrictive temperature. That this is the case is demonstrated by a radioimmune assay that detects material that cross-reacts with antibody directed against

purified T4 DNA polymerase. Despite the fact that little or no T4 DNA polymerase activity appeared at 43°, cross-reacting material (CRM) for the antipolymerase antibody could be detected with the radioimmune assay (Table 1). In fact, as much or more CRM was made at the nonpermissive temperature as active enzyme made at the permissive temperature. It was also found (Table 2) that the level of CRM increased after infection, reaching a maximum at about 20 min postinfection. This direct measurement correlates well with the kinetics of formation of the precursor inferred from the T4 DNA polymerase activity ultimately achieved upon step down after various periods at 43° (Swartz *et al.*, 1972). Even those temperature-sensitive mutants, L91 and L141, that did not form active polymerase, nevertheless did produce substantial amounts of CRM (data not shown).

*The 3' → 5' exonuclease activity of T4 DNA polymerase undergoes activation.* Despite the lack of polymerase activity at the restrictive temperature, it was possible that the CRM made at 43° might retain the 3' → 5' exonuclease activity of T4 DNA polymerase. However, as shown in Fig. 6, the 3' → 5' exonuclease activity was absent at 43° and only appeared when the infected




FIG. 5. Time course of T4 DNA polymerase appearance after infection of JT1200 by various temperature-sensitive mutants in gene 43. Separate cultures (200 ml) of JT1200 growing at 43° in HT-broth were infected at an  $A_{595}$  of 0.8 with the mutant T4 phages indicated at an m.o.i. of 5 each. The experiment was performed as described for Fig. 1. The triangles (*ts* L97, *ts* S9 and *ts* L141) show the level of activation achieved in separate experiments with a less detailed time course. The closed circles (*ts* L88 and *ts* L141) represent experiments in which the initial temperature was 45°.

TABLE 1  
CRM PRODUCED BY T4 MUTANTS IN GENES 43 AND 44<sup>a</sup>

T4 phage	Gene	Temperature of infection (degrees)	T4 DNA polymerase	
			Activity	CRM (units/mg protein)
<i>ts</i> L56 <i>am</i> N82	43, 44	43	0.5	19.2
<i>ts</i> L56 <i>am</i> N82	43, 44	30	6.3	5.9
<i>am</i> N82	44	37	6.9	8.7
<i>am</i> 4309	43	37	0.3	0.2

<sup>a</sup> Cultures of *E. coli* B growing at the temperatures indicated were infected at an  $A_{595}$  of 0.8 with the T4 phages listed at an m.o.i. of 5. After 15 min, T4 DNA polymerase and CRM activities were assayed at 30° in Brij lysates of the infected cultures as described under Materials and Methods.

TABLE 2  
TIME COURSE OF APPEARANCE OF CRM<sup>a</sup>

Time after infection (min)	CRM (units/mg protein)
0	< 1
10	10
20	13.4
40	10.5
60	8.7

<sup>a</sup> A culture of *E. coli* B growing at 43° was infected at an  $A_{595}$  of 0.8 with T4 *ts* L56 *am*N82 at an m.o.i. of 5. At the times indicated, portions of the culture were withdrawn and assayed for CRM as described under Materials and Methods.

cells were shifted to 30°, exactly as is found for the polymerase activity.

*Energy for activation is derived from "high energy" phosphate.* Several experiments suggest that the energy necessary for activation can be derived from either oxidative or substrate-level phosphorylations, and thus, it probably represents the need for some high energy phosphate compound (or compounds). First, as demonstrated in Fig. 7, at no time after lowering the temperature did activation escape the need for metabolic energy. Apparently, energy-yielding metabolism must operate continuously to provide a large enough pool of the required component for the activation process to go to completion. Second, arsenate was an effective inhibitor of the




FIG. 6. Time course of appearance of polymerase and 3' → 5' exonuclease activities of T4 DNA polymerase. A culture (200 ml) of JT1200 growing at 43° in HT-broth was infected at an  $A_{595}$  of 0.8 with T4 *ts* L56 at an m.o.i. of 5. The experiment was performed as described for Fig. 1, except that assays for exonuclease activity were carried out in the presence of 100  $\mu$ l of anti-*E. coli* exonuclease I serum. The 3' → 5' exonuclease activity of T4 DNA polymerase is that exonuclease activity present in each of the extracts that was inhibited by 25  $\mu$ l of anti-T4 DNA polymerase serum.

activation process, with half-maximal effect at 2–3 mM (Fig. 8). This concentration of arsenate had no effect on the activity of the purified polymerase. Addition of arsenate has been shown to lower drastically the intracellular pools of ATP, phosphoenolpyruvate, and other “energy-rich” phosphate compounds (Klein and Boyer, 1972; Berger, 1973). Third, activation occurred normally in a mutant of *E. coli*, designated *uncA*<sup>-</sup>, in which the Ca<sup>2+</sup>, Mg<sup>2+</sup>-stimulated ATPase associated with oxidative phosphorylation is defective and which therefore must obtain its ATP from the substrate-level phosphorylations of glycolysis (Fig. 9). In addition, activation in the *uncA*<sup>-</sup> strain proceeded essentially normally in the presence of 2,4-dinitrophenol, but was effectively inhibited by arsenate (data not shown). Fourth, *de novo* deoxyribonucleotide biosynthesis was not required for activation since the addition of 50 mM hydroxyurea, an inhibitor of both the cellular and the phage-induced ribonucleotide reductase (Neuhard, 1967; Warner and Hobbs, 1969), either at 5 min after infection or at the time of temperature

downshift, did not block subsequent activation (data not shown).

*Covalent modification of the protein does not occur upon activation.* Because of the evidence *in vivo* for the participation of high-energy phosphate compounds in the activation process, the possible involvement of covalent modifications that have been described in other systems (Walsh *et al.*, 1968; Kingdon *et al.*, 1967; Honjo *et al.*, 1968) was investigated. Our results suggest that activation does not involve phosphorylation, adenylation, or adenosine diphosphoribosylation of the enzyme. Purified preparations of active T4 DNA polymerase contained no detectable phosphorus by direct chemical analysis under conditions where the requisite number of phosphate residues were found for a known phosphoprotein (pepsinogen). Furthermore, there is no shoulder at 260 nm in the ultraviolet adsorption spectrum of purified T4 DNA polymerase (Goulian *et al.*, 1968). In addition, when infected cells prelabeled




FIG. 7. Effect of 2,4-dinitrophenol on the appearance of T4 DNA polymerase. A 350-ml culture of *E. coli* B growing at 43° in H-broth was infected at an  $A_{595}$  of 0.8 with T4 *ts* L56 at an m.o.i. of 5. The experiment was performed as described in Fig. 1 except that at the times indicated, 30-ml samples were withdrawn and their incubation continued in the presence of a final concentration of 1 mM 2,4-dinitrophenol, while the remainder of the culture was incubated at 30° in the absence of 2,4-dinitrophenol.




FIG. 8. Effect of various concentrations of arsenate on the time course of T4 DNA polymerase appearance. A culture (400 ml) of *E. coli* B growing at 43° in HT-broth was infected at an  $A_{595}$  of 0.8 with T4 *ts* L56 at an m.o.i. of 5. The experiment was performed as described for Fig. 2, except that at the time of temperature shift, arsenate was added to different portions of the culture. ■, 100 mM arsenate; ●, 50 mM arsenate; □, 20 mM arsenate; ▲, 10 mM arsenate; ○, 5 mM arsenate; △, 2 mM arsenate; ◇, 1 mM arsenate; ●, no arsenate.

with  $[^{32}\text{P}]\text{PO}_4^{3-}$  were used in a typical activation experiment, and the proteins extracted from the cells at various times after infection were radioautographed after sodium dodecyl sulfate-polyacrylamide slab gel electrophoresis, no radioactivity was found associated with the band corresponding to T4 DNA polymerase. Thus, it must be concluded that the enzyme is neither phosphorylated, adenylylated, nor adenosine diphosphoribosylated, either transiently or permanently, during the process of activation. Also, unlike several structural proteins of the phage (Celis *et al.*, 1973; Goldstein and Champe, 1974; Bachrach and Benchetrit, 1974; Poglazov and Levschenko, 1974; Georgopoulos *et al.*, 1972), T4 DNA polymerase does not appear to be generated from a larger molecular weight precursor polypeptide. When cells were pulse-labeled with  $[^{35}\text{S}]\text{SO}_4^{2-}$  at various times after infection in a typical activation experiment, a labeled species that comigrated on dodecyl sulfate-polyacrylamide slab gels with authentic T4 DNA polymerase appeared at about 5 min after infection. This band was the largest molecular weight species present

and at no time before or after temperature downshift was its mobility different from that of the purified enzyme within the resolution of this technique. In fact, when the CRM for anti-T4 DNA polymerase was precipitated from extracts of cells infected at the restrictive temperature, a discrete band was found with a mobility indistinguishable from that of  $^{125}\text{I}$ -labeled purified T4 DNA polymerase marker.

#### DISCUSSION

It is well established that the primary sequence of a protein determines its tertiary structure (Epstein *et al.*, 1963), however, it has been stressed by many investigators (Alpers and Paulus, 1971; Teipel




FIG. 9. Time course of T4 DNA polymerase appearance in a mutant defective in oxidative phosphorylation. Separate cultures (600 ml) of *E. coli* AN180 and AN120 were grown at 43° in HT-broth to an  $A_{595}$  of 0.4. The cells were chilled and collected by brief centrifugation at 8000 rpm. They were then resuspended in one-twentieth their original volume of phage absorption buffer (0.1 M NaCl, 2 mM  $\text{MgCl}_2$ , 10  $\mu\text{g}/\text{ml}$  tryptophan, and 20 mM triethanolamine hydrochloride, pH 7.5) and were infected at 0° with T4 *ts* L56 at an m.o.i. of 5. After allowing adsorption to occur for 10 min, the cells were rapidly diluted into 110 ml of HT-broth prewarmed to 43°. The rest of the experiment was as described in Fig. 1.

and Koshland, 1971; Frieden, 1970; Hatfield and Burns, 1970) that other factors may influence the ultimate conformation that a protein assumes and may dictate whether or not that state of the protein will be functional. Therefore, it might be expected that mutations will exist that will affect not only the activity of enzymes, but also their formation.

We have previously found that after infection of *E. coli* at 43° with the temperature-sensitive T4 mutant *tsL56*, an inactive polymerase was synthesized that was rapidly converted to active enzyme on shift to a permissive temperature (30°). Although the conversion occurred in the absence of protein synthesis, it was prevented by inhibitors of energy-yielding metabolism, such as sodium azide, sodium cyanide, or 2,4 dinitrophenol.

In the present study, we have demonstrated that activation of T4 DNA polymerase is not an allele-specific effect; a variety of mutations in gene 43, which confer temperature-sensitivity on the activity of T4 DNA polymerase, also affect the formation of a functional enzyme. Furthermore, at least in the case of *tsL56*, the activation affects both polymerase and exonuclease activities.

We have also shown that during infection at the nonpermissive temperature, cross-reacting material for T4 DNA polymerase specific antibody is made. Two to three times as much CRM accumulates at the nonpermissive temperature as active enzyme formed at the permissive temperature (Tables 1 and 2). These findings can be explained most simply as an increased accessibility of the antigenic sites on the unfolded precursor.

Our *in vivo* experiments strongly suggest the participation of "high energy" phosphate in the activation process. On the other hand, direct comparisons of the inactive precursor and the functional enzyme indicate that the energy-dependent activation does not involve phosphorylation, adenylation, or adenosine diphosphoribosylation of the protein, although such covalent modifications have been detected in other phage systems (Rahmsdorf *et al.*, 1974; Horvitz, 1974a, 1974b; Goff, 1974; Seifert *et al.*, 1971). Also, despite the

existence in *E. coli* of an energy-dependent system responsible for the degradation of "missense" proteins (Prouty and Goldberg, 1972; Goldberg, 1972), activation does not appear to involve proteolytic cleavage of the inactive precursor. Of course, covalent modifications of the protein during activation can be ruled out definitely only by purification of the inactive precursor and the active enzyme, followed by peptide fingerprinting and sequencing of the two proteins.

Although it seemed reasonable to suppose that the energy-dependent activation may be related to insertion of the protein into a replication complex and that mutations that affect interaction of T4 DNA polymerase with other T4 proteins may interfere with its ability to form the complex, mutations that should have a drastic effect on its ability to form the replication complex did not prevent the appearance of active T4 DNA polymerase.

The gene 43 *amber* mutant B22 (Fig. 4), which lacks 20% of the carboxyl-terminal portion of the protein, retains its 3' → 5' exonuclease despite complete loss of polymerase activity (Nossal and Hershfield, 1972). However, temperature-sensitive mutations in the carboxyl-terminal region of the protein (*tsL88*, L91, and L141) prevent activation of both nuclease and polymerase activities. Taken together, these results suggest that an intact carboxyl region of the protein is somehow necessary for activation, and that its absence is preferable to a defect in this region for the production of the enzymatically active molecule.

Our first attempts to prepare an *in vitro* system capable of generating active T4 DNA polymerase met with only limited success (Swartz *et al.*, 1972). The involvement of transient intermediates or labile components may have been responsible, in part, for the difficulty, especially since the assay for activation requires the completion of a multistep process. Indeed, the extent of activation that can be achieved in the various cell-free systems that we have attempted to date is only 10% or so of the levels obtainable *in vivo*.<sup>7</sup> Moreover, while

<sup>7</sup> J. Thorner and I. R. Lehman, unpublished experiments.

this limited *in vitro* activation occurs most rapidly at 30°, nevertheless, it will proceed upon storage of extracts at 4°. Despite these problems, preparation and fractionation of an *in vitro* system seems the best approach toward elucidating the mechanism of the activation process.

## REFERENCES

- ALLEN, E. F., ALBRECHT, I., and DRAKE, J. W. (1970). Properties of bacteriophage T4 mutants defective in DNA polymerase. *Genetics* **65**, 187-200.
- ALPERS, J. B., and PAULUS, H. (1971). Allosteric preconditioning: Role of allosteric ligands in promoting the maturation of enzymes. *Nature (London)* **233**, 478-479.
- APOSHIAN, H. V., and KORNBERG, A. (1962). Enzymatic synthesis of deoxyribonucleic acid. IX. The polymerase formed after T2 bacteriophage infection of *Escherichia coli*: A new enzyme. *J. Biol. Chem.* **237**, 519-525.
- BACHRACH, U., and BENCHETRIT, L. (1974). Studies on phage internal proteins. II. Cleavage of a precursor of internal proteins during the morphogenesis of bacteriophage T4. *Virology* **59**, 51-58.
- BARRY, J., and ALBERTS, B. (1972). *In vitro* complementation as an assay for new proteins required for bacteriophage T4 DNA replication: Purification of the complex specified by T4 genes 44 and 62. *Proc. Nat. Acad. Sci. USA* **69**, 2717-2721.
- BERGER, E. A. (1973). Different mechanisms of energy coupling for the active transport of proline and glutamine in *Escherichia coli*. *Proc. Nat. Acad. Sci. USA* **70**, 1514-1518.
- BRUTLAG, D., and KORNBERG, A. (1972). Enzymatic synthesis of deoxyribonucleic acid. XXXVI. A proofreading function for the 3' → 5' exonuclease activity in deoxyribonucleic acid polymerase. *J. Biol. Chem.* **247**, 241-248.
- BUTLIN, J. D., COX, G. B., and GIBSON, F. (1971). Oxidative phosphorylation in *Escherichia coli* K-12: Mutations affecting magnesium ion- or calcium ion-stimulated adenosine triphosphatase. *Biochem. J.* **124**, 75-81.
- CELIS, J. E., SMITH, J. D., and BRENNER, S. (1973). Correlation between genetic and translational maps of Gene 23 in bacteriophage T4. *Nature New Biol.* **241**, 130-132.
- CHEN, P. S., TORIBARA, T. Y., and WARNER, H. (1956). Microdetermination of phosphorus. *Analyt. Chem.* **28**, 1756-1758.
- DEWAARD, A., PAUL, A. V., and LEHMAN, I. R. (1965). The structural gene for deoxyribonucleic acid polymerase in bacteriophages T4 and T5. *Proc. Nat. Acad. Sci. USA* **54**, 1241-1248.
- EPSTEIN, C. J., GOLDBERGER, R. F., and ANFINSEN, C. B. (1963). The genetic control of tertiary protein structure: Studies with model systems. *Cold Spring Harbor Symp. Quant. Biol.* **28**, 439-449.
- EPSTEIN, R. H., BOLLE, A., STEINBERG, C. M., KELLENBERGER, E., BOY DE LA TOUR, E., CHEVALLEY, R., EDGAR, R. S., SUSMAN, M., DENHART, G. H., and LIELAUSIS, A. (1963). Physiological studies of conditional lethal mutants of bacteriophage T4D. *Cold Spring Harbor Symp. Quant. Biol.* **28**, 375-394.
- FREIDEN, C. (1970). Kinetic aspects of regulation of metabolic processes. The hysteretic enzyme concept. *J. Biol. Chem.* **245**, 5788-5799.
- GEORGIOPOULOS, C. P., HENDRIX, R. W., KAISER, A. D., and WOOD, W. B. (1972). Role of the host cell in bacteriophage morphogenesis: Effects of a bacterial mutation on T4 head assembly. *Nature New Biol.* **239**, 38-41.
- GOFF, C. (1974). Chemical structure of a modification of the *Escherichia coli* ribonucleic acid polymerase polypeptides induced by bacteriophage T4 infection. *J. Biol. Chem.* **249**, 6181-6190.
- GOLDBERG, A. L. (1972). Degradation of abnormal proteins in *Escherichia coli*. *Proc. Nat. Acad. Sci. USA* **69**, 422-426.
- GOLDSTEIN, J., and CHAMPE, S. P. (1974). T4-induced activity required for specific cleavage of a bacteriophage protein *in vitro*. *J. Virol.* **13**, 419-427.
- GOULIAN, M., LUCAS, Z. J., and KORNBERG, A. (1968). Enzymatic synthesis of deoxyribonucleic acid. XXV. Purification and properties of deoxyribonucleic acid polymerase induced by infection with phage T4. *J. Biol. Chem.* **243**, 627-638.
- HALL, Z., and LEHMAN, I. R. (1968). An *in vitro* transversion by a mutationally altered T4-induced DNA polymerase. *J. Molec. Biol.* **36**, 321-333.
- HATFIELD, G. W., and BURNS, R. O. (1970). Ligand-induced maturation of threonine deaminase. *Science* **167**, 75-76.
- HAYES, W. (1968). "The Genetics of Bacteria and Their Viruses." John Wiley and Sons, New York.
- HONJO, T., NISHIZUKA, Y., HAYAISHI, O., and KATO, I. (1968). Diphtheria toxin-dependent adenosine diphosphate ribosylation of aminoacyl transferase II and inhibition of protein synthesis. *J. Biol. Chem.* **243**, 3553-3555.
- HORVITZ, H. R. (1974a). Control by bacteriophage T4 of two sequential phosphorylations of the  $\alpha$  subunits of *Escherichia coli* RNA polymerase. *J. Molec. Biol.* **90**, 727-738.
- HORVITZ, H. R. (1974b). Bacteriophage T4 mutants deficient in alteration and modification of the *Escherichia coli* RNA polymerase. *J. Molec. Biol.* **90**, 739-750.
- HUANG, W. M., and BUCHANAN, J. M. (1974). Synergistic interactions of T4 early proteins concerned with their binding the DNA. *Proc. Nat. Acad. Sci. USA* **71**, 2226-2230.
- HUANG, W. M., and LEHMAN, I. R. (1972a). On the exonuclease activity of phage T4 deoxyribonucleic acid polymerase. *J. Biol. Chem.* **247**, 3139-3146.

- HUANG, W. M., and LEHMAN, I. R. (1972b). On the direction of translation of the T4 deoxyribonucleic acid polymerase gene *in vivo*. *J. Biol. Chem.* **247**, 7663-7667.
- HUBERMAN, J. A., KORNBERG, A., and ALBERTS, B. M. (1971). Stimulation of T4 bacteriophage DNA polymerase by the protein product of T4 gene 32. *J. Molec. Biol.* **62**, 39-52.
- KINGDON, H. S., SHAPIRO, B. M., and STADTMAN, E. R. (1967). Regulation of glutamine synthetase. VIII. ATP:glutamine synthetase adenylyltransferase, an enzyme that catalyzes alterations in the regulatory properties of glutamine synthetase. *Proc. Nat. Acad. Sci. USA* **58**, 1703-1710.
- KLEIN, W. L., and BOYER, P. D. (1972). Energization of active transport. *J. Biol. Chem.* **247**, 7257-7265.
- KORNBERG, T., and GEFTER, M. L. (1971). Purification and DNA synthesis in cell-free extracts: Properties of DNA polymerase. II. *Proc. Nat. Acad. Sci. USA* **68**, 761-764.
- KORNBERG, T., and GEFTER, M. L. (1972). Deoxyribonucleic acid synthesis in cell-free extracts. IV. Purification and catalytic properties of deoxyribonucleic acid polymerase. III. *J. Biol. Chem.* **247**, 5369-5375.
- LEHMAN, I. R., and CHIEN, J. R. (1973). Persistence of DNA polymerase I and its 5' → 3' exonuclease activity in *PolA* mutants of *Escherichia coli* K-12. *J. Biol. Chem.* **248**, 7717-7723.
- LOWRY, O. H., ROSEBROUGH, N. J., FARR, A. L., and RANDALL, R. J. (1951). Protein measurement with the folin phenol reagent. *J. Biol. Chem.* **193**, 265-275.
- MILLER, J. H. (1972). "Experiments in Molecular Genetics," Cold Spring Harbor Laboratory, Cold Spring Harbor, New York.
- NEUHARD, J. (1967). Studies on the acid-soluble nucleotide pool in *Escherichia coli*. IV. Effects of hydroxyurea. *Biochim. Biophys. Acta* **145**, 1-6.
- NOSSAL, N. G., and HERSHFIELD, M. S. (1971). Nuclease activity in a fragment of bacteriophage T4 deoxyribonucleic acid polymerase induced by the amber mutant B22. *J. Biol. Chem.* **246**, 5414-5426.
- POGLAZOV, B. F., and LEVSHENKO, M. T. (1974). Bacteriophage T4D-induced proteinase. *J. Molec. Biol.* **84**, 463-467.
- PROUTY, W. F., and GOLDBERG, A. L. (1972). Fate of abnormal proteins in *E. coli*. Accumulation in intracellular granules before catabolism. *Nature New Biol.* **240**, 147-150.
- RAHMSDORF, H. J., PAL, S. H., PONTA, H., HERRLICH, P., ROSKOSKI, R., SCHWEIGER, M., and STUDIER, F. W. (1974). Protein kinase induction in *Escherichia coli* by bacteriophage T7. *Proc. Nat. Acad. Sci. USA* **71**, 586-589.
- SEIFERT, W., RABUSSEY, D., and ZILLIG, W. (1971). On the chemical nature of alteration and modification of DNA-dependent RNA polymerase after T4 infection. *FEBS Lett.* **16**, 175-178.
- STEINBERG, C. M., and EDGAR, R. S. (1967). A critical test of a current theory of genetic recombination in bacteriophage. *Genetics* **47**, 187-208.
- SWARTZ, M. N., NAKAMURA, H., and LEHMAN, I. R. (1972). Activation of a defective DNA polymerase in a temperature-sensitive mutant of bacteriophage T4. *Virology* **47**, 338-353.
- TAYLOR, A. L., and TROTTER, C. D. (1972). Linkage map of *Escherichia coli* strain K-12. *Bact. Rev.* **36**, 504-524.
- TEIPEL, J. W., and KOSHLAND, D. E., JR. (1971). Kinetic aspects of conformational changes in proteins. I. Rate of regain of enzyme activity from denatured proteins. *Biochemistry* **10**, 792-798.
- WALSH, D. A., PERKINS, J. P., and KREBS, E. G. (1968). An adenosine 3',5'-monophosphate-dependent protein kinase from rabbit skeletal muscle. *J. Biol. Chem.* **243**, 3763-3774.
- WARNER, H. R., and HOBBS, M. D. (1967). Incorporation of uracil-<sup>14</sup>C into nucleic acids in *Escherichia coli* infected with T4 and T4 amber mutants. *Virology* **33**, 376-384.
- WARNER, H. R., and HOBBS, M. D. (1969). Effect of hydroxyurea on replication of bacteriophage T4 in *Escherichia coli*. *J. Virol.* **3**, 331-336.